Effects of bark beetle and drought induced tree mortality on wildfire severity in the Sierra Nevada

Rebecca Wayman¹ and Hugh Safford^{1,2}

1. University of California, Davis 2. USDA Forest Service Pacific Southwest Region

Study Question: Does recent tree mortality influence wildfire severity in an historically frequent fire forest?



Rough fire, 8/9/2015. National Park Service.

Rough fire, 9/2015. USDA Forest Service.

Vegetation and fuel dynamics following severe pine mortality (theoretical)



Disturbance Characteristics and Metrics

Pre-fire mortality conditions		Fire conditions		
Outbreak phase (time since outbreak)	Outbreak metric	Fire metric	Fire severity type	Historic fire regime
Red (1-3 yrs)	Area affected	Likelihood	Canopy	High frequency, low/mod severity
Gray (3-10 yrs)	Severity	Severity	Surface	Low/mod frequency, high severity
Old/tree fall (11-20 yrs)		Intensity		

Study area



2015 Rough Fire ~60,000 ha (~150,000 ac)

2016 Cedar Fire ~12,000 ha (~30,000 ac)

Study area: 2016 Cedar Fire



Study area: 2015 Rough Fire





Rough fire approaching the Kings River, USDA Forest Service 2015.

Does recent tree mortality influence wildfire severity in an historically frequent fire forest?

If yes, does the relationship vary under different weather or stand structural conditions?

Mortality and fire severity metrics



Predictor = mortality conditions *immediately pre-fire*

Mortality and fire severity metrics



Response = mortality conditions *immediately post-fire*

Field methods



Fir engraver(*Scolytus ventralis*) and gallery on white fir (*Abies concolor*). A. Brackett 2017.

- One-year post-fire field assessment of fire severity
- Determination of the cause and timing of tree mortality

Field methods



Dead at the time of fire:

- 1. Old phase
- 2. Gray phase
- 3. Red phase

Live at the time of fire:

- 4. Killed by fire
- 5. Killed by insects post-fire
- 6. Green attack
- 7. No attack

Mountain pine beetle (*Dendroctonus ponderosae*) and wood borer (*Melanophila* sp.) galleries on sugar pine (*Pinus lambertiana*) killed by beetles prior to wildfire. A. Brackett 2017.

Results: Influential Variables on Cedar Fire Severity



Influential variables as identified by random forests analysis. Variables to the right of the red line are considered influential to the percent of plot basal area killed by fire.

Results: Influential Variables on Rough Fire Severity



Influential variables as identified by random forests analysis. Variables to the right of the red line are considered influential to the percent of plot basal area killed by fire.

Pre-Fire Mortality Matters During More Severe Weather, Cedar Fire



Pre-Fire Mortality Matters at Higher Basal Area, Rough Fire



Pre-Fire Mortality is More Influential than Stand Density

Density and Pre-fire Mortality Interaction



Generalized Additive Model

Full Model = Best Model (R² adj. = 0.595):Total density (live + dead)Pre-fire dead tree density*Stand basal area*Topographic relative moisture index*Relative humidity on burn date*Total density x pre-fire dead density*Stand basal area x pre-fire dead density*

*significant at p<=0.001

Acknowledgements



Data collection on the 2016 Cedar fire, June 2017.

Thanks to Beverly Bulaon and crew members Amanda Brackett, Amanda Cooke, Katherine Kangas, Tessa Putz.



Data collection on the 2015 Rough fire, June 2016.

Thank You

contacts: rbwayman@ucdavis.edu hdsafford@ucdavis.edu

Background photo credits (Cedar fire): USDA Forest Service 2016