#### University of California Agriculture and Natural Resources Making a Difference for California



### UCCE/DWR Weekly Crop Water Use Report

#### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET<sub>C</sub>) 05/30/25 through 06/05/25

Crops (Leafout Date)	3	#148 Merced			#39 Parlier				#258 Lemon Cove		
	05/30 - 06/05	Accum'd	06/06 - 06/12		05/30 - 06/05	Accum'd	06/06 - 06/12		05/30 - 06/05	Accum'd	06/06 - 06/12
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc
Almonds (3/1) *	1.73	14.00	1.81		1.87	14.74	1.81		1.81	13.63	1.81
Pistachio (4/25) * **	1.80	5.70	2.03		1.99	6.28	2.02		1.92	5.90	2.03
Citrus (2/1)	1.29	13.69	1.40		1.41	14.48	1.39		1.37	13.44	1.40
Raisin Grapes (4/14) (11 ft. row spacing)	1.19	5.38	1.30		1.33	5.84	1.30		1.28	5.46	1.30
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.22	5.26	1.34		1.34	5.71	1.33		1.30	5.36	1.34
Walnuts (4/14)	1.39	6.33	1.54		1.52	6.86	1.54		1.47	6.46	1.54
Stone Fruit (3/8)	1.39	8.51	1.50		1.51	9.15	1.50		1.46	8.51	1.50
Past 7 days precipitation (inches)		0.00				0.00		_		0.00	
Accumulated precipitation (inches) (1/1/2025)		0.00				5.35				4.44	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

DACE STEELER	A DDT TED 337 A TED	O THE INCIDED	A D THOMED DO	D DEDUCTED OF 1
PAST WEEKLY	APPLIED WATE	R IN INCHES.	ADJUSTED FO	R EFFICIENCY 1

Crops		#148 Merce	d			#39 Parlier				#258 Lemon	n Cove	
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.7	2.3	2.0	1.8	2.9	2.5	2.2	2.0	2.8	2.4	2.1	1.9
Pistachio (4/25)	2.8	2.4	2.1	1.9	3.1	2.7	2.3	2.1	3.0	2.6	2.3	2.0
Citrus (2/1)	2.0	1.7	1.5	1.4	2.2	1.9	1.7	1.5	2.1	1.8	1.6	1.4
Raisin Grapes (4/14) (11 ft. row spacing)	As	sume all gra	pe	1.3	Assume all grape 1.4			1.4	A	1.3		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.3	irrigation type is drip			1.4	irrigation type is drip			1.4
Walnuts (4/14)	2.1	1.9	1.6	1.5	2.3	2.0	1.8	1.6	2.3	2.0	1.7	1.5
Stone Fruit (3/8)	2.1	1.9	1.6	1.5	2.3	2.0	1.8	1.6	2.2	1.9	1.7	1.5

<sup>1</sup> The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

#### PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#148 Merced					#39 Parlier			#258 Lemon Cove								
Almonds 115 Trees/A	638	543	472	425	685	590	519	472	661	567	496	449					
Pistachio 106 Trees/A	698 598 523		473	772	673	573	523	747	648	573	498						
Citrus 110 Trees/A	494	420	370	346	543	469	420	370	518	444	395	346					
Raisin Grapes 566 Vines/A	Assume all grape			62	Assume all grape 67			67	Assume all grape			62					
Winegrapes 622 Vines/A	irrig	ation type is	drip	57	irrigation type is drip 61			61	irrigation type is drip			61					
Walnuts 76 Trees/A	750	679	572	536	822	715	643	572	822	715	607	536					
Stonefruit 172 Trees/A	332 300 253		237	363	316	284	253	347	300	268	237						
For further information concerning all counties receiving this report, contact	the Fresno Co	. Farm Advis	or's office at	(559) 241-75	26.	for further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.											

# University of California Agriculture and Natural Resources

Making a Difference for California



## UCCE/DWR Weekly Crop Water Use Report

#### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or  $ET_C$ ) 05/30/25 through 06/05/25

Crops (Leafout Date)	#	#124 Panoche			#	2 Five Point	ts	#15 Stratford			
	05/30- 06/05	Accum'd	06/06- 06/12		05/30- 06/05	Accum'd	06/06- 06/12	05/30- 06/05	Accum'd	06/06- 06/12	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	1.85	14.56	2.01		1.90	14.95	2.02	2.07	16.04	2.04	
Pistachio (4/25) * **	1.98	6.36	2.23		2.03	6.53	2.24	2.20	6.99	2.26	
Citrus (2/1)	1.40	14.88	1.53		1.44	15.20	1.54	1.57	16.35	1.56	
Raisin Grapes (4/14) (11 ft. row spacing)	1.31	5.93	1.44		1.35	6.11	1.45	1.46	6.56	1.46	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.32	5.81	1.48		1.35	5.93	1.48	1.46	6.39	1.50	
Walnuts (4/14)	1.52	7.02	1.74		1.55	7.13	1.75	1.69	7.64	1.77	
Stone Fruit (3/8)	1.49	9.24	1.70		1.54	9.36	1.71	1.66	9.99	1.73	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2025)		2.31				3.13			2.72		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHE	S, ADJUSTED FOR EFFICIENCY 1
#124 Damasha	#2 E: D-:4-

Crops	#124 Panoche					#2 Five Poi	nts		#15 Stratford			
System Efficiency>>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.8	2.5	2.2	1.9	2.9	2.5	2.2	2.0	3.2	2.8	2.4	2.2
Pistachio (4/25)	3.0	2.6	2.3	2.1	3.1	2.7	2.4	2.1	3.4	2.9	2.6	2.3
Citrus (2/1)	2.2	1.9	1.6	1.5	2.2	1.9	1.7	1.5	2.4	2.1	1.8	1.7
Raisin Grapes (4/14) (11 ft. row spacing)	As	sume all gra	pe	1.4	Assume all grape 1.4			1.4	Assume all grape			1.5
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.4	irrigation type is drip		drip	1.4	irrigation type is drip		drip	1.5
Walnuts (4/14)	2.3	2.0	1.8	1.6	2.4	2.1	1.8	1.6	2.6	2.3	2.0	1.8
Stone Fruit (3/8)	2.3	2.0	1.8	1.6	2.4	2.1	1.8	1.6	2.6	2.2	2.0	1.7

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

DACT WEEKI V	APPLIED WATER	INCALLONDED	TDEE OD VINE

Crops		#124 Panoc	he			#2 Five Poi	nts			#15 Stratfo	rd	
Almonds 115 Trees/A	661	590	519	449	685	590	519	472	756	661	567	519
Pistachio 106 Trees/A	747	648	573	523	772	673	598	523	847	722	648	573
Citrus 110 Trees/A	543	469	395	370	543	469	420	370	592	518	444	420
Raisin Grapes 566 Vines/A	As	ssume all gra	ipe	67	Assume all grape 67			67	Assume all grape			72
Winegrapes 622 Vines/A	irrig	ation type is	drip	61	irrigation type is drip			61	irrigation type is drip			65
Walnuts 76 Trees/A	822	715	643	572	857	750	643	572	929	822	715	643
Stonefruit 172 Trees/A	363	316	284	253	379	332	284	253	410	347	316	268
	(1 F C	г 41'	1 00 4	(550) 241 7	1537							

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

<sup>\*</sup> Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.