University of California Agriculture and Natural Resources Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 06/27/25 through 07/03/25

Crops (Leafout Date)	;	#148 Merced			#39 Parlier			#258 Lemon Cove		
	06/27 - 07/03	Accum'd	07/04 - 07/10		06/27 - 07/03	Accum'd	07/04 - 07/10	06/27 - 07/03	Accum'd	07/04 - 07/10
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc
Almonds (3/1) *	2.09	21.81	2.10		2.19	23.16	2.03	2.14	21.60	2.06
Pistachio (4/25) * **	2.17	14.10	2.17		2.27	15.28	2.10	2.22	14.46	2.13
Citrus (2/1)	1.47	19.38	1.47		1.57	20.64	1.40	1.52	19.24	1.43
Raisin Grapes (4/14) (11 ft. row spacing)	1.41	10.80	1.43		1.45	11.60	1.37	1.42	10.91	1.39
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.58	11.18	1.61		1.65	12.08	1.55	1.61	11.38	1.57
Walnuts (4/14)	1.84	13.07	1.96		1.94	14.19	1.89	1.89	13.36	1.92
Stone Fruit (3/8)	1.82	15.13	1.91		1.92	16.36	1.84	1.87	15.31	1.87
Past 7 days precipitation (inches)		0.00				0.00			0.00	
Accumulated precipitation (inches) (1/1/2025)		0.00				5.35			4.44	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

Due to July 4th holiday, data for 07/03/2025 used an estimated Eto value

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1

THE TWEEKET MITERED WATER BY BUILDING TOR ELITICIES OF													
Crops		#148 Merce	d			#39 Parlier			#258 Lemon Cove				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	3.2	2.8	2.5	2.2	3.4	2.9	2.6	2.3	3.3	2.9	2.5	2.3	
Pistachio (4/25)	3.3	2.9	2.6	2.3	3.5	3.0	2.7	2.4	3.4	3.0	2.6	2.3	
Citrus (2/1)	2.3	2.0	1.7	1.5	2.4	2.1	1.8	1.7	2.3	2.0	1.8	1.6	
Raisin Grapes (4/14) (11 ft. row spacing)	As	ssume all gra	pe	1.5	Assume all grape 1.5			1.5	A	1.5			
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.7	irrigation type is drip			1.7	irrigation type is drip			1.7	
Walnuts (4/14)	2.8	2.5	2.2	1.9	3.0	2.6	2.3	2.0	2.9	2.5	2.2	2.0	
Stone Fruit (3/8)	2.8	2.4	2.1	1.9	3.0	2.6	2.3	2.0	2.9	2.5	2.2	2.0	

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops		#148 Merce		#39 Parlier			#258 Lemon Cove							
Almonds 115 Trees/A	756	661	590	519	803	685	614	543	779	685	590	543		
Pistachio 106 Trees/A	822	722	648	573	872	747	673	598	847	747	648	573		
Citrus 110 Trees/A	568	494	420	370	592	518	444	420	568	494	444	395		
Raisin Grapes 566 Vines/A	Assume all grape			72	Assume all grape 72			72	Assume all grape			72		
Winegrapes 622 Vines/A	irrig	gation type is	drip	74	irrigation type is drip 74			74	irrigation type is drip			74		
Walnuts 76 Trees/A	1000	893	786	679	1072	929	822	715	1036	893	786	715		
Stonefruit 172 Trees/A	442	379	332	300	474	410	363	316	458	395	347	316		

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

University of California Agriculture and Natural Resources Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 06/27/25 through 07/03/25

Crops (Leafout Date)	#	#124 Panoche			#2 Five Points			#15 Stratford			
	06/27- 07/03	Accum'd	07/04- 07/10		06/27- 07/03	Accum'd	07/04- 07/10	06/27- 07/03	Accum'd	07/04- 07/10	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	2.13	22.92	2.17		2.25	23.98	2.23	2.35	25.27	2.24	
Pistachio (4/25) * **	2.21	15.31	2.24		2.33	16.12	2.30	2.42	16.77	2.31	
Citrus (2/1)	1.51	20.98	1.54		1.61	21.79	1.60	1.66	23.04	1.61	
Raisin Grapes (4/14) (11 ft. row spacing)	1.43	11.65	1.43		1.50	12.30	1.48	1.58	12.89	1.49	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.59	12.06	1.67		1.69	12.69	1.72	1.76	13.29	1.73	
Walnuts (4/14)	1.88	14.29	2.03		2.00	14.99	2.09	2.09	15.69	2.10	
Stone Fruit (3/8)	1.86	16.39	1.98		1.98	17.09	2.06	2.07	17.89	2.08	
Past 7 days precipitation (inches)		0.00		_		0.00		 	0.00		
Accumulated precipitation (inches) (1/1/2025)		2.31				3.13			2.72		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

Due to July 4th holiday, data for 07/03/2025 used an estimated Eto value

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIE											
Crops		#124 Panoch	ıe	#2 Five Points							
			0 - 0 /				0 -0 /				

Crops			#2 Five Poi	nts								
System Efficiency>>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.3	2.8	2.5	2.2	3.5	3.0	2.6	2.4	3.6	3.1	2.8	2.5
Pistachio (4/25)	3.4	2.9	2.6	2.3	3.6	3.1	2.7	2.5	3.7	3.2	2.8	2.5
Citrus (2/1)	2.3	2.0	1.8	1.6	2.5	2.1	1.9	1.7	2.6	2.2	2.0	1.7
Raisin Grapes (4/14) (11 ft. row spacing)	A	ssume all gra	pe	1.5	Assume all grape 1.6			1.6	As	1.7		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	gation type is	drip	1.7	irrigation type is drip			1.8	irrigation type is drip			1.9
Walnuts (4/14)	2.9	2.5	2.2	2.0	3.1	2.7	2.4	2.1	3.2	2.8	2.5	2.2
Stone Fruit (3/8)	2.9	2.5	2.2	2.0	3.0	2.6	2.3	2.1	3.2	2.8	2.4	2.2

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PACT WEEKI V	APPI IFN WATER	IN GALLON PER	TREE OR VINE

Crops			#2 Five Poi	nts		#15 Stratford						
Almonds 115 Trees/A	779	661	590	519	826	708	614	567	850	732	661	590
Pistachio 106 Trees/A	847	722	648	573	897	772	673	623	922	797	698	623
Citrus 110 Trees/A	568	494	444	395	617	518	469	420	642	543	494	420
Raisin Grapes 566 Vines/A	Assume all grape			72	Assume all grape 77			77	Assume all grape			82
Winegrapes 622 Vines/A	irrig	ation type is	drip	74	irrigation type is drip 79			79	irrigation type is drip			83
Walnuts 76 Trees/A	1036	893	786	715	1108	965	857	750	1143	1000	893	786
Stonefruit 172 Trees/A	458	395	347	316	474	410	363	332	505	442	379	347

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.