2.7

2.4

University of California Agriculture and Natural Resources Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

3.0

2.6

2.3

2.1

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 07/25/25 through 07/31/25

Crops (Leafout Date)	1	#148 Merced			#39 Parlier			#258 Lemon Cove		
	07/25 - 07/31	Accum'd	08/01 - 08/07		07/25 - 07/31	Accum'd	08/01 - 08/07	07/25 - 07/31	Accum'd	08/01 - 08/07
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc
Almonds (3/1) *	2.00	30.01	1.95		1.98	31.45	1.88	1.92	29.61	1.84
Pistachio (4/25) * **	2.07	22.58	2.02		2.05	23.85	1.94	1.96	22.72	1.86
Citrus (2/1)	1.40	25.09	1.39		1.39	26.47	1.32	1.34	24.84	1.28
Raisin Grapes (4/14) (11 ft. row spacing)	1.34	16.35	1.30		1.33	17.19	1.24	1.28	16.27	1.21
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.52	17.43	1.48		1.50	18.41	1.42	1.45	17.48	1.39
Walnuts (4/14)	2.14	21.35	2.09		2.12	22.56	2.01	2.03	21.42	1.93
Stone Fruit (3/8)	2.07	23.11	2.02		2.05	24.45	1.94	1.96	23.07	1.86
Past 7 days precipitation (inches)		0.00				0.00			0.00	
Accumulated precipitation (inches) (1/1/2025)		0.00				5.35			4.44	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

Stone Fruit (3/8)

* Estimates are for orchard floor conditions where vegetation is managed by some combination of sarip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY													
Crops	#148 Merced				#39 Parlier				#258 Lemon Cove				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	3.1	2.7	2.4	2.1	3.0	2.6	2.3	2.1	3.0	2.6	2.3	2.0	
Pistachio (4/25)	3.2	2.8	2.4	2.2	3.2	2.7	2.4	2.2	3.0	2.6	2.3	2.1	
Citrus (2/1)	2.2	1.9	1.6	1.5	2.1	1.9	1.6	1.5	2.1	1.8	1.6	1.4	
Raisin Grapes (4/14) (11 ft. row spacing)	A	ssume all gra	ipe	1.4	Assume all grape			1.4	Assume all grape			1.3	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.6	irrigation type is drip			1.6	irrigation type is drip		1.5		
Walnuts (4/14)	3.3	2.9	2.5	2.3	3.3	2.8	2.5	2.2	3.1	2.7	2.4	2.1	

DACT WEEKLY ADDITED WATED IN INCHES ADDICTED FOR EFFICIENCY.

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE													
Crops	#148 Merced				#39 Parlier				#258 Lemon Cove				
Almonds 115 Trees/A	732	638	567	496	708	614	543	496	708	614	543	472	
Pistachio 106 Trees/A	797	698	598	548	797	673	598	548	747	648	573	523	
Citrus 110 Trees/A	543	469	395	370	518	469	395	370	518	444	395	346	
Raisin Grapes 566 Vines/A	As	ssume all gra	pe	67	Assume all grape 67			67	Assume all grape			62	
Winegrapes 622 Vines/A	irrig	ation type is	drip	70	irrigation type is drip		70	irrigation type is drip		65			
Walnuts 76 Trees/A	1179	1036	893	822	1179	1000	893	786	1108	965	857	750	
Stonefruit 172 Trees/A	505	442	379	347	505	426	379	347	474	410	363	332	
For further information concerning all counties receiving this report, contact	the Fresno Co	. Farm Adviso	or's office at	(559) 241-75	26.					•			

UCCE/DWR Weekly Crop Water Use Report





WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 07/25/25 through 07/31/25

Crops (Leafout Date)		#124 Panoch				2 Five Point	to	#15 Stratford				
Crops (Leafout Date)											<u> </u>	
	07/25- 07/31	Accum'd	08/01- 08/07		07/25- 07/31	Accum'd	08/01- 08/07	07/25- 07/31	Accum'd	08/01- 08/07		
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated		
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc		
Almonds (3/1)*	1.96	31.10	1.95		2.08	32.74	2.10	2.11	34.13	2.10		
Pistachio (4/25) * **	2.03	23.77	2.02		2.15	25.14	2.17	2.18	25.88	2.17		
Citrus (2/1)	1.38	26.72	1.39		1.45	27.94	1.47	1.48	29.26	1.47		
Raisin Grapes (4/14) (11 ft. row spacing)	1.31	17.14	1.30		1.39	18.15	1.43	1.42	18.81	1.43		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.49	18.27	1.48		1.59	19.34	1.61	1.62	20.01	1.61		
Walnuts (4/14)	2.10	22.55	2.09		2.22	23.81	2.24	2.25	24.60	2.24		
Stone Fruit (3/8)	2.03	24.36	2.02		2.15	25.70	2.17	2.18	26.60	2.13		
Past 7 days precipitation (inches)		0.00				0.00			0.00			
Accumulated precipitation (inches) (1/1/2025)		2.31				3.13			2.72			

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUST	TED FOR EFFICIENCY 1
---	----------------------

TAST WEEKLT ATTEMED WATER IN INCHES, ADJUSTED FOR EFFICIENCY													
Crops	#124 Panoche				#2 Five Points				#15 Stratford				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	3.0	2.6	2.3	2.1	3.2	2.8	2.4	2.2	3.2	2.8	2.5	2.2	
Pistachio (4/25)	3.1	2.7	2.4	2.1	3.3	2.9	2.5	2.3	3.4	2.9	2.6	2.3	
Citrus (2/1)	2.1	1.8	1.6	1.5	2.2	1.9	1.7	1.5	2.3	2.0	1.7	1.6	
Raisin Grapes (4/14) (11 ft. row spacing)	As	ssume all gra	pe	1.4	Assume all grape 1.5			1.5	Assume all grape			1.5	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.6	irrigation type is drip			1.7	irrigation type is drip		1.7		
Walnuts (4/14)	3.2	2.8	2.5	2.2	3.4	3.0	2.6	2.3	3.5	3.0	2.6	2.4	
Stone Fruit (3/8)	3.1	2.7	2.4	2.1	3.3	2.9	2.5	2.3	3.4	2.9	2.6	2.3	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#124 Panoche					#2 Five Poi	nts		#15 Stratford			
Almonds 115 Trees/A	708 614 543			496	756	661	567	519	756	661	590	519
Pistachio 106 Trees/A	772	673	598	523	822	722	623	573	847	722	648	573
Citrus 110 Trees/A	518	444	395	370	543	469	420	370	568	494	420	395
Raisin Grapes 566 Vines/A	Assume all grape			67	Assume all grape 72			72	Assume all grape			72
Winegrapes 622 Vines/A	irrig	ation type is	drip	70	irrigation type is drip 74			74	irrigation type is drip			74
Walnuts 76 Trees/A	1143	1000	893	786	1215	1072	929	822	1251	1072	929	857
Stonefruit 172 Trees/A	489	426	379	332	521	458	395	363	537	458	410	363
For further information concerning all counties receiving this report, contact	For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.											

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.